

Adapting practice: Infection risk assessment and mitigation policy

This document provides a written record of the heightened infection control measures that this clinic has put into place to ensure the safety of all staff and patients during COVID-19 and thereafter.

We have assessed our practice for risks outlined and put in additional processes as detailed below • 21.09.2022 – replacing previous risk assessment of 03.02.2022. Instigated with relaxation of all COVID 19 restrictions Undertaken a risk assessment **Heightened cleaning** Bathroom will be cleaned after every usage where possible regimes Carpeted areas will be steam cleaned daily Increased protection Removal of towels and fabric plinth covering and pillowcases from bathroom and clinic room Contactless and BACS payment preferred measures Put in place Appointments every 45 minutes, allowing airing and cleaning between patients distancing measures • Chairs placed 2m apart in waiting room Staff training We are aware of correct handwashing technique best practice, Putting on & removal of PPE safely We compiled updated clinic policies and infection measures together Providing pre-• All patients will have telephone pre-screening call screening telephone calls and advice remotely where prudent Document last updated: 21.09.2022

	Description of risk	Mitigating action	When introduced
Pre-screening for risk before public/patients visit the clinic	Exposure to COVID with inherent risk of developing the illness ourselves, spread to family members, patients and any person entering the house	 The following will be applied to patients and their chaperone if applicable - screening for additional respiratory symptoms or conditions e.g. hay fever, asthmas etc Screen to see if a member of their household had/has symptoms of COVID-19 Have they been in contact with someone with suspected/confirmed COVID-19 in last 7days? 	21.09.2022
Protecting members of staff	If shielding a vulnerable member of our respective households there is risk of exposure to COVID-19	Neither Lynette or Clover fall into this category	
Confirmed cases of COVID 19 amongst staff or patients?	Risk of exposure to osteopath treating that patient, patients entering the clinic, members of their households	 If the patient experiences symptoms within 2/3 days of visiting the clinic, any staff with direct contact to that individual should request a lateral flow test 	
Travel to and from the clinic	Public transport/taxi – risk of exposure	We shall ask patients to stay in their car or wait until they see the previous patient leave	
Entering and exiting the building	Exposure to virus through close proximity to other patients	 asking patients not to arrive early or late for their appointment patients arriving early will be asked to wait in their car or if arrival on foot, taxi or public transport and in inclement weather observe social distancing in the waiting room patients will be expected use the hand sanitiser provided in the porch before entering the building 	21.09.2022

	Description of risk	Mitigating action	When introduced
Reception and common areas	contact with other patients	 ask patients to turn up promptly at their appointment time to reduce time in the waiting area encourage contactless payment and BACS instead of cash, on making their appointment 	
Social/physical distancing measures in place	proximity and risk of contamination	 appointments every 45 minutes to allow airing and cleaning between patients Chairs will be place 2m apart in the waiting room and clinic room 	
Face to face consultations (in-clinic room)	Exposure to virus through closer than 2m contact, aerosol risk through techniques	 Modifying treatment techniques to avoid unnecessary close proximity One parent/guardian only with visits for children No additional family members except if requested as a chaperone Pre-screen chaperones as with patients prior to appointment 	•

Table 2b Hygiene mea We have assessed the		f risk in our practice and put in place the following heightened hygiene mea Mitigating action	Sures When introduced
Increased sanitisation and cleaning	Exposure to virus and spread to other families and patients	 Sanitising Clinic rooms - plinths, desk, door handles, touch points, equipment chairs - between each patient Doors and door handles, chairs, taps, card machines etc. in other areas Use of at least 60% alcohol sanitisers/wipes Remove unnecessary linen/use plastic pillowcases that can be cleaned between patients Provide single use blankets which can washed at high temperature in the colder weather Using disposable paper towels in the bathroom Decluttering the clinic rooms and waiting area of unnecessary items 	21.09.2022

	Description of	Mitigating action	When
	risk		introduced
Aeration of rooms	As above	 Keeping treatment door open after each patient Aeration of common/reception areas by keeping windows open in good weather. In poor weather opening doors and windows after every patient and at the end of every working session i.e. morning or afternoon 	
Staff hand hygiene measures	As above	 Bare below the elbow/hand washing before and after patients with soap and water for at least 20 seconds, including forearms/use of hand sanitiser gel/ use of gloves 	
Respiratory and cough hygiene	As above	 cough hygiene measures for osteopaths and patients - 'Catch it, bin it, kill it' Provision of disposable, single-use tissues waste bins (lined and foot-operated) Hand hygiene facilities available for patients, visitors, and osteopaths 	
Cleaning regimes	As above	 Cleaning common areas at the end of each working session. Frequent cleaning of the bathroom 	

Table 3. Personal Protective Equipment: Detail here your policy for use and disposal of PPE		
Clinicians will wear the following PPE	 Fluid-resistant surgical masks worn where considered prudent and on request from patient Eye protection, e.g. if there is a risk of droplet transmission or fluids entering eyes Use of visors where patient is deaf and communication is difficult without ability to lip read due to face mask 	
When will PPE be replaced	 When potentially contaminated, damaged, damp, or difficult to breathe through At the end of a session – morning or afternoon or up to 4 hours duration 	
Reception staff will wear the following PPE	Not applicable	
Patients will be asked to wear the following PPE	Wearing of masks by patients is optional	

PPE disposal	Double-plastic bagged, away from other household/garden waste, and then incinerated

Table 4. Communicating with patients: Detail here how you will advise patients of measures that we have taken to ensure their safety and the policies that have been put in place in our clinic		
Publishing your updated clinic policy	 Printed and displayed on hall table Available on Paignton Osteopaths website 	
Information on how you have adapted practice to mitigate risk	 Available for viewing on Paignton Osteopaths website Advice monitored and updated following Government guidance 	
Pre-appointment screening calls	 when making the appointment with a request to contact the clinic if anything changes in accordance with screening call criteria A clinician will call. 	
Information for patients displayed in the clinic	 Door notices advising anyone with symptoms not to enter the building. Notices on other public health measures e.g. hand washing/sanitising Providing patients contact for more information if needed 	